

FOR CONFUSED FLOUR BEETLES, SAWTOOTHED GRAIN BEETLES, GRANARY WEEVILS, RICE WEEVILS, RED FLOUR BEETLES, GRAIN MITES: Spray into cracks and crevices, processing and handling equipment, around cartons or containers and other areas where these insects tend to congregate. Spray 30 seconds for each 100 square feet of surface. Keep area closed for 45 minutes to 1 hour after treatment. Repeat application as necessary.

FOR FLEAS: Spray sleeping quarters, bedding, floor and floor covering where pets are kept. Repeat as necessary. For best results, use with a recommended pet spray on pets.

To kill Cockroaches, Spiders, Silverfish, Wasps, Hornets, Crickets – Spray the insect directly when possible and spray all probable hiding places. Repeat as necessary.

To kill Flies, Mosquitoes, Gnats, Small Flying Moths – Close all windows and doors. Direct spray upward, spraying for 2-3 seconds for 1,000 cubic feet of space. Do not remain in treated areas. Ventilate after treatment.

VOLUMETRIC TREATMENT FOR INACCESSIBLE VOIDS (Attics, False Ceilings, Wall and Equipment Voids): For crack and crevice application, calculate void volume in cubic feet. Treat 20 seconds per 1,000 cubic feet. Repeat as necessary.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Do not store near heat or open flame. Store in cool dry area away from children.

DISPOSAL: Do not Puncture or Incinerate!

If empty: Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions.



PRECAUTIONARY STATEMENTS Hazards to Humans & Domestic Animals CAUTION

Harmful if swallowed or absorbed through skin. Avoid breathing of vapors or spray mist. Avoid contact with skin and eyes. In case of contact, immediately flush eyes or skin with plenty of water. Obtain medical attention if irritation persists. Remove pets, birds and cover fish aquariums before spraying.

PHYSICAL OR CHEMICAL HAZARDS

Flammable. Contents under pressure. Keep away from heat, sparks and open flames. Do not puncture or incinerate container. Exposure to temperatures above 130°F may cause bursting. Do not spray on plastic, painted or varnished surfaces or directly into electronic equipment such as radios, TVs, computers, etc.

FIRST AID

If Swallowed: Call a poison control center or doctor for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. **If Inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. **If on Skin or Clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice. **If In Eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **HOT LINE NUMBER** – Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information, call (985) 878-6751, Monday through Friday, 8:00 am to 4:30 pm CST. After 4:30pm call your poison control center. For product information call (203) 597-1812, Monday-Friday, 8:00 am to 5:00 pm EST.



INSTANT KNOCKDOWN

CV-80

FARM & DAIRY
For Insect Control

0.50% Pyrethrins
for Quick KILL

Provides Rapid Control of Barn Pests

KILLS & REPELS: Flies, Mosquitos, Small Flying Moths, Gnats, Fleas, Barn Flies, Deer Flies, Stable Flies, Horn Flies, Horse Flies and House Flies

FOR USE IN:

Beef Cattle Operations, Dairy Farms, (Including Milk House, Milk Parlor, Lofing Sheds and Holding Lot), Hog Operations, Barns, Stables, Farms, Animal Quarters, Milkrooms and Poultry Houses



CONTAINS NO CFCs OR OTHER OZONE DEPLETING SUBSTANCES.
FEDERAL REGULATIONS PROHIBIT CFC PROPPELLANTS IN AEROSOLS.

ACTIVE INGREDIENTS:

Pyrethrins.....0.50%
*Piperonyl Butoxide, TECHNICAL.....4.00%
INERT INGREDIENTS.....95.50%
*Equivalent to 3.2% (butylcarbityl) (6-propylpiperonyl) ether and 0.8% related compounds. 100.00%

KEEP OUT OF REACH OF CHILDREN
CAUTION

See Back Panel for Additional Precautionary Statements.

EPA Reg. No. 9444-175 EPA Est. No. 9444-LA-1

NET WT 17 oz (482 g)

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Note, when using this product in an enclosed area, such as a milkroom, barn or stable, remove animals before spraying.

DAIRY FARM USE: Milking Parlor and Milkroom: Close all windows and doors. Direct spray upwards, spraying for 2-3 seconds for 1,000 cubic feet of space. Do not remain in treated areas. Ventilate after treatment. Cover milking utensils and milk before spraying to prevent contamination from dead and falling insects.

ANIMAL USE: Thoroughly spray entire animal from approximately 2 feet distance. Do not spray directly toward animal's eyes. Spot treat withers, shoulders, and back where saliva accumulates from head tossing.

BEEF CATTLE OPERATIONS: In barns, close all windows and doors. Spray at a rate of 2-3 seconds per 1,000 cubic feet of space. Do not remain in treated area. After 15 minutes, ventilate the area. Repeat application if necessary.

ANIMAL USE: For animal application, thoroughly spray entire animal from approximately 2 feet distance. Do not spray directly towards animal's face or eyes. Repeat application daily or as necessary.

STANCHION BARN USE: Walk behind animals and direct spray over backs, allowing 5 seconds per animal. For most effective results, apply each morning.

HORSE BARN OR STABLES: In horse and pony barns, close all windows and doors. Spray at the rate of 2-3 seconds per 1,000 cubic feet of space. Do not remain in treated area. After 15 minutes, ventilate the area. Repeat the application as necessary.

ANIMAL USE: For animal application, lightly mist over back of horses and ponies from a distance of 2 feet. Do not spray directly toward the animal's face or eyes. Repeat application daily or as necessary.

HOG OPERATIONS: In hog houses, close all doors and windows. Spray at the rate of 2-3 seconds per 1,000 cubic feet of space. Do not remain in treated area. After 15 minutes, ventilate the area. Repeat the application as necessary.

SWINE USE: For animal application, direct spray over back of hogs and spray for 5 seconds per hog. Do not spray directly toward the animal's face or eyes. Repeat the application daily or as necessary.

POULTRY OPERATIONS: In poultry houses, close all windows and doors. Spray roosts, walls, nests and cages thoroughly. Spray at the rate of 2-3 seconds per 1,000 cubic square feet of space. Do not remain in treated areas. After 15 minutes, ventilate the area. Treat daily or as necessary.

FLYING INSECTS: Flies, Mosquitoes, Small Flying Moths, Gnats – Calculate cubic feet of space to be treated. Close all doors and windows. Spray in sweeping motion to all parts of room. Apply for 2 to 3 seconds per 1,000 cubic feet. Sweep up and destroy fallen insects. Repeat treatment as necessary.

CRAWLING INSECTS: Cockroaches, Ants, Spiders, Silverfish – General Volumetric Treatment: Open cabinets and doors in area to be treated. Shut off air conditioners, fans, and close doors and windows before spraying. Spray for 20 seconds per 1,000 cubic feet of space. Direct spray toward areas suspected of harboring insects. Spray into hiding places such as cracks and crevices, behind sinks, cabinets, along baseboards and floors, around drains and plumbing, hitting insects with spray whenever possible. Keep area closed for 15 minutes. Open and ventilate before reoccupying. Repeat treatment as necessary.

FOR COCKROACHES: For best results, spray directly on cockroaches and thoroughly spray all parts of room suspected of harboring these pests. Special attention should be paid to cracks, hidden surfaces, under sinks, behind stove and refrigerators, food storage areas, and wherever these pests are suspected of hiding. Also, spray around door and window frames. Repeat as necessary.

FOR SPIDERS AND SILVERFISH: For control of these insects, frequent and repeated spraying of baseboards, windows, doorframes, and storage areas should be made. For best results, spray directly on insects.

FOR CRICKETS: Spray thoroughly and frequently in storage areas and especially in other damp areas. For best results, spray directly on insects.

FOR ANTS: Spray ant trails, nests, and points of entry. For initial clean-up, apply for 10-15 seconds per 100 square feet of floor area. Then periodically spray as a space treatment for 3-5 seconds per 100 cubic feet of space.

FOR INDIANMEAL MOTHS, ANGOUIMOIS GRAIN MOTHS: Close doors and windows. Direct spray into all parts of the room, especially around stored food product containers, pallets, and darkened areas. Spray 7 to 10 seconds for each 1,000 cubic feet of space. Keep area closed for 15 minutes following treatment. Ventilate after treatment. Repeat application weekly or as needed.

Part #34-7310CVR



Manufactured by
WATERBURY COMPANIES, INC.

P.O. Box 1812 • Waterbury, CT 06722
985-878-6751 • Made in U.S.A.

